

COLLISION INDUCED ELECTRONIC TRANSITIONS (CIET) IN NESTED SINGLET EXCITED STATES OF N₂: A TWO-COLOR PUMP-PROBE REMPI STUDY^a

MURTHY S. GUDIPATI^b, MARSHALL L. GINTER, *Institute of Physical Sciences and Technology, University of Maryland, College Park, MD 20742*; RICHARD A. COPELAND, *Molecular Physics Laboratory, SRI International, 333 Ravenswood Ave, Menlo Park, CA 94025*.

Collision induced electronic transitions (CIET) occur between two or more different electronic states that lie energetically close to one another. If these electronic states are associated with different vibrational anharmonicities, dipole moment or polarizabilities, then near-resonance conditions can occur between some of the vibronic levels of these nested electronic states in different surroundings. Well-known examples of such processes include radiative and non-radiative relaxation among the triplet states of CO.

In addition, CIET plays a very important role in the processes that occur in earth's upper atmosphere, where airglow and aurora occur through excitation of atomic and molecular nitrogen and oxygen species by electrons from solar wind, photons from the sun or atom recombination.

In order to get accurate rate constants for CIET occurring in N₂ between the excited singlet states $a^1\Pi_g$ and $a'^1\Sigma_u^-$, we have carried out two-color pump-probe REMPI experiments. The pump laser is used to populate initially $v = 1$ level of the $a^1\Pi_g$ state through a two-photon excitation. The probe laser ionizes N₂ from the time-evolved electronic states through a one-photon resonance to highly excited singlet states of N₂. Our results show that direct collisional relaxation from $v = 1$ to $v = 0$ of the $a^1\Pi_g$ state is far less efficient (less than 10%) than CIET to generate $v = 1$ of the $a'^1\Sigma_u^-$ state. Further details of our investigations and their relevance to atmospheric processes will be presented and discussed.

^aThis project is supported by NSF Grant No. 9910914

^bVisiting Scientist at SRI International (on leave from University of Cologne, Germany)

Time required: 15 min

Session in which paper is recommended for presentation: 2