

Near-Infrared Emission of Highly Excited CO – A More Sensitive Method to Probe the Inter Stellar Matter??

Murthy S. Gudipati^{1,2}

¹*Institute of Physical Sciences and Technology, University of Maryland,
College Park, MD 20742*

²*Visiting Scientist at Molecular Physics Laboratory, SRI International,
333 Ravenswood Avenue, Menlo Park, CA 94025*

Carbon monoxide is one of the major constituents of stellar and interstellar media. The near infrared emission of CO due to the ground-state vibrational transitions at 4.67 and 2.3 μm are used as markers to study the evolution and ages of several stars [1]. The vacuum ultraviolet (VUV) features of CO have been observed with *IUE* followed by *HST* using *GHR*S-Spectrograph [2]. On the other hand, the spectroscopic features of CO in the near infrared region (NIR) between 0.9 and 2 μm involving the electronically excited states have not yet been explored in astrophysical observations.

There are four triplet electronic states of CO that lie energetically close to each other like nested ladders between the ground-state ($X^1\Sigma^+$) and the first optically allowed excited singlet state ($A^1\Pi$). These triplet states with increasing energy are: $a^3\Pi$, $a'^3\Sigma^+$, $d^3\Delta$ and $e^3\Sigma^-$, respectively. The a -state is a metastable state with a lifetime of several ms; 94 ms in the gas-phase, 90 ms in Ne, and 7.2 ms in Ar matrices [3]. Hence, the transitions involving this a -state and the other three upper states, covering the wavelength region between UV and NIR, should be of importance for astrophysical studies of CO-abundant objects.

In our experimental work, we have observed that the band-maxima of the NIR emission of CO are highly sensitive to the environment in which CO exists. Due to the fact that the above mentioned triplet states are associated with opposing dipole-moments, these states also shift energetically in opposing directions with increasing polarizability of the medium.

Based on the large shifts observed in the positions of these NIR emission bands, we have proposed a few years ago [4], that a combination of astrophysical observations and laboratory experiments in the NIR region on CO would be a high-sensitive method to probe stellar and interstellar objects containing CO and are exposed to VUV radiation. Further details of the photophysical processes will be presented.

References:

- [1] Greene, T.P., Lada, C.J., *ApJ*, **461**, 345 (1996)
- [2] Lambert, D.L., Sheffer, Y., Gilliland, R.L., Federman, S.R., *ApJ*, **420**, 756 (1994)
- [3] Fournier, J., Mohammed, H.H., Deson, J., Vermeil, C., Schamps, J., *J. Chem. Phys.*, **73**,6039 (1980)
- [4] Gudipati, M. S., Kalb, M., *Astron. Astrophys.*, **329**, 375 (1998)

Acknowledgments:

This work was carried out at the Synchrotron Radiation Facility in Berlin. The Author is also affiliated with the University of Cologne, Germany.